

**TWO NEW TRIASSIC EUTOMOCERID AMMONOID CEPHALOPODS
(AMMONOIDEA: CERATITIDAE) FROM SOUTHEASTERN
PENINSULAR THAILAND**

Kittichai Tongtherm¹, Jaruwat Nabhitabhata^{2*} and Surangkana Tuanapaya³

¹*Graduate School of Life and Environmental Sciences, University of Tsukuba,
Tsukuba, Ibaraki 305-8572, Japan*

²*Excellence Centre for Biodiversity of Peninsular Thailand (CBIPT), Faculty of Science,
Prince of Songkla University, Hatyai, Songkhla 90112, Thailand*

³*Reference Collection, Phuket Marine Biological Center, Muang District, Phuket 83000, Thailand*

**Corresponding author: jaruwat.n5@gmail.com*

ABSTRACT: Two ceratitid ammonoids of *Eutomoceras* are described as new species based on characters in the internal mold of the phragmocone. The specimens were collected from Khao Ok-Thalu, Chaiburi Formation, Triassic strata in Changwat Phatthalung, southeastern Peninsular Thailand. The two new species share characters of bifurcate ribs, absence of marginal furrows on the ventral keel and round ventrolateral shoulders. The umbilicus is wide with the absence of knots on the shoulder in *E. paladeji* sp. nov., but it is narrower with the presence of knots in *E. wantanae* sp. nov. The new species bring the number of described *Eutomoceras* species to seven. The identification key to species of *Eutomoceras* is proposed herein.

Keywords: new species, Triassic Strata, ceratitid ammonoids, *Eutomoceras*, Peninsular Thailand
